
**Effect of Drinking Water Supplementation With Extract
Fermented *Tamarindus Indica* By *Saccharomyces Spp* On Egg
Production, Feed Digestibility And N-NH₃ Content In Excreta
Laying Hens**

Mateus Da Cruz De Carvalho*, Ida bagus Gaga Partama, I Gusti
Nyoman Gde Bidura*** & Ni Wayan Siti******

&*Animal Science Department, Agricultural Faculty of National University, Timor Lorosa'e, East
Timor*

&*Doctor Programs of Animal Sci., Faculty of AniM. Sci., Udayana University, Indonesia*

ABSTRACT

*The present experiment was conducted to determine effects of drinking water supplementation with extract fermented *Tamarindus indica* by *Saccharomyces spp* (Tamarin-Pro) on egg production, feed digestibility, and N-NH₃ in excreta of laying hens up to 54 weeks old. A total of 240 laying hens with an average initial body weight of 1674.91±18.52 g were randomly allotted into four drinking water treatments. The drinking water treatments included: drinking with 2% Tamarin-Pro extract (A); drinking water with 4% Tamarin-Pro extract (B); drinking water with 6% Tamarin-Pro extract (C); and drinking water as a control (D), respectively. Result showed that the Tamarin-Pro drinking water treatment had improved feed digestibility compared with control treatments ($p < .05$). The egg mass in Tamarin-Pro treatments B and D were greater than that in control treatment ($p < .05$). Moreover, Tamarin-Pro extract decreased significantly ($p < .05$) N-NH₃ in excreta of birds. In conclusion, fermented *Tamarindus indica* by *Saccharomyces spp* (Tamarin-Pro) extract supplementation improved egg mass and nutrient digestibility in laying hens and decreased N-NH₃ in excreta of hens*

KEY WORDS: *Tamarindus indica*, water extract, probiotics, ammonia, feed digestibility

INTRODUCTIONS

The trend of using phytochemical compounds from herbs in feed and drinking water has increased over the past two decades. The results of the study showed that the use of antibiotics in excessive feed would intensify the potential risks of increasing resistance in human pathogens. Bacterial resistance and antibiotic residues in animal products cause increasing concern in using antibiotics as a growth driver and ultimately result in a ban on the use of antibiotics in feed in most developed countries (Gheisar and Kim, 2018)

The herbal leaf fermentation process can increase the efficacy of the treatment of active ingredients and reduce the anti-nutritional effects on herbal leaves, thereby increasing the growth of poultry and the value of the benefits of herbal leaves (Ahmed *et al.*, 2016). Tanin is known to have a bitter or astringent taste which reduces palatability and hence food intake (Jansman, 1993). The fermentation process can reduce the concentration of tannic acid, which can reduce the adverse effects of tannin found in natural herbs on feed intake.

Supplementation of herbal plant extracts can improve taste and feed consumption, stimulate animal appetite, and then increase feed intake (Frankic *et al.*, 2009). As a result, increased feed intake can contribute to increased productivity of poultry. Different results regarding the effects of herbal extract additives on livestock production performance were reported in the study of Gheisar and Kim (2018) and Ahmed *et al.* (2017). Similar results were reported by Ahmed *et al.* (2016) that supplementation of herbal leaves (pomegranate, Gingko biloba and licorice) in natural or fermented forms, turned out to have no effect on growth, but reduced feed consumption and increased feed efficiency compared to controls (without herbal leaf supplementation).

The phytobiotic antioxidant function of herbal leaves is positively influencing food stability, improving egg quality, and prolonging the storage time of eggs (eggs are easily damaged if stored too long). In general, some results of the study show the positive effects of herbal phytochemical compounds on livestock performance (Gheisar and Kim, 2018; Lei *et al.*, 2018; Bidura *et al.*, 2017) and probiotics (Bidura *et al.*, 2019; Tufarelli *et al.*, 2017; Phuoc and Jamikorn, 2017; Mountzouris *et al.*, 2010). These conflicting results regarding growth performance responses to natural herbs, fermentation, or extractions can be caused by different herb species, concentrations of herbs, and methods of processing herbs (Windisch *et al.*, 2008; Hashemi and Davoodi, 2011; Embuscado 2015).

Recently, in Indonesia there has been a ban on the use of antibiotics, so probiotics and phytochemicals of herbal leaves have been suggested as the most beneficial alternative for livestock because of their beneficial effects (Abdelqader *et al.*, 2013). *Tamarindus indica* leaves have long been used by the community as herbal leaves with their phytochemical content, such as alkaloid, steroid, terfenoid, fenolik, flavonoid, dan tanin (Carvalho, 2019). Likewise, the yeast *Saccharomyces sp* has long been used as a "tape" fermentation inoculant, and can act as probiotics to improve feed digestibility and poultry performance, and reduce ammonia gas content in cages (Bidura *et al.*, 2012; 2019 and Ezema and Eze, 2015). Hasan *et al.* (2016) and Hasanuddin *et al.* (2017; Zhen *et al.*, 2019) reported that that feed fermentation generally improves the bacterial ecology of the digestive tract and the immune response in chicks, therefore, becomes a new model for future strategies to control chicken disease.

It is interesting to study the use of medicinal properties of phytochemicals combined with probiotics. Phytochemical and probiotic compounds, both show antimicrobial effects, antioxidants, anti-inflammatory, boost growth and improve feed efficiency. Therefore this study was conducted to examine the use of *Tamarindus indica* leaf extract fermented by probiotics *Saccharomyces spp* (*Tamarin-Pro*) on drinking water to egg production, feed digestibility, and ammonia levels in excreta of laying hens

MATERIAL AND METHODS

Experimental design, animals, housing and diets

A total of 240 laying hens with an average initial body weight of 1674.91 ± 18.52 g were randomly allotted into four drinking water treatments. The drinking water treatments included: drinking water with 2% *Tamarin-Pro* extract (A); drinking water with 4% *Tamarin-*

Pro extract (B); drinking water with 6% *Tamarin-Pro* extract (C); and drinking water as a control (D), respectively. All hens were housed in an environmentally controlled room with forced ventilation. Each pen was equipped with a nipple drinker and a varalon feeder. Laying hens were provided with free access to drinking water and feed throughout experimental period. All chickens were given commercial feed specific for laying hens containing 2.750 kcal/kg of Metabolizable Energy (ME); 18% of CP; 3.5% of Ca; and available phosphor of 0.45%.

The laying hens were randomly divided into four groups: A: drinking water with administration of 2% *Tamarin-Pro* water extract; B: with the administration of 4% *Tamarin-Pro* water extract; C: with the administration of 6% *Tamarin-Pro* water extract; and D: drinking water without administration of *Tamarin-Pro* water extract, respectively. Each treatment consisted of five replicate pens with 10 birds were randomly assigned to each pen at 150×70×45 cm (length×width×height). Each experimental diet was in mash form and the birds had free access to feed and water throughout the experiment.

Preparation of natural and fermented herbs

The leaves of *Tamarindus indica* are washed, dried in air and powdered. For solid fermentation, *Tamarindus indica* leaf powder is mixed with *Saccharomyces spp* at 3.7×10^7 colony forming units (cfu)/g. After that, the mixture is immersed in distilled water to maintain a moisture concentration of 40%. The hydrated herb is then fermented at 37°C. After fermentation for 72 hours, the fermented sample is mixed with distilled water 1: 1 (v: v) and it is meserated for 24 hours. Then filtered to get *Tamarindus* leaf water extract which has been fermented with *Saccharomyces spp* as a probiotic source. This water extract is then referred to as *Tamarin-Pro* extract.

Live performance.

Continuous lighting and access to feed and water was provided throughout the experiment. The birds were weighed at the commencement (40 weeks of age) and the end (56 weeks of age) of the experiment. Eggs were collected daily and egg production was expressed on a hen-day basis (% hens day). Individual egg weights were recorded then used to calculate mean egg weight for all experimental period. The total egg mass was calculated by multiplying egg weights by egg production. Feed intake was measured on a cage basis (hen) every week. Daily feed intake per bird was calculated on a cage total feed intake basis for the entire experimental period and for the number of days in all the period. Feed conversion ratio (kg of feed/kg of eggs mass) for the all period was calculated on a cage basis from egg production, egg weight, and feed consumption.

Retention and excretion of nutrients.

In order to determine the nutrient digestibility (dry matter and organic matter digestibility) value of the diet: The amount of diet used was 100 g, this amount as based on preliminary assays with laying hens consumption of ration. All the birds were deprived of feed for 24 h to ensure that their alimentary canals were empty from feed residues. They were then force-fed with the specific amount of diets (all treatments). Stainless steel funnel with 40 cm stem was used in *force feeding technique* (Bidura *et al.*, 2019). Water was available *ad libitum* during the experimental period. The total excreta were collected in plastic trays. The excreta samples were frozen, allowed coming to equilibrium with the atmospheric moisture, weighed,

and ground through a 1 mm sieve. Samples of excreta and diets were subjected to appropriate analysis to determine dry matter (DM) and organic matter (OM), respectively. Dry matter (DM) and organic matter (OM), and ash determinations were done according to the Association of Official Analytical Chemists (2005). All assays were conducted in triplicate.

Determination of N-NH₃ concentration in excreta

The determination of N-NH₃ levels in excreta was carried out using the Conway dish diffusion as follows: 1 ml of supernatant sample on the left side of the Conway dish, 1 ml of saturated Na₂CO₃ solution on the right bulkhead, 1 ml of H₃BO₃ 2% indicated BCG+MR on the middle cup, then the vaseline conway cup is closed tightly, shake it slowly, until the supernatant with Na₂CO₃ solution is completely mixed, then left for 24 hours at room temperature, then titrate using a 0.005N H₂SO₄ solution, until the end point titration. N-NH₃ levels can be calculated as follows: mM N-NH₃ = (titration volume x N H₂SO₄ x 1,000)

Statistical analysis

All data were analyzed with ANOVA to determine the differences among treatments. If differences were found, then further analysis was performed with Duncan's multiple range test.

RESULTS

The results showed that initial weight, final body weight, number of eggs, feed consumption, feed conversion ratio, digestibility of dry matter, organic matter, and ammonia levels in excreta in group A (drinking water with 2% extract of *Tamarin-Pro*), B (drinking water with 4% extract of *Tamarin-Pro*), C (drinking water with 6% extract of *Tamarin-Pro*), and D (drinking water without *Tamarin-Pro* extract as control), respectively were presented as in Table 1. The total number of eggs and feed consumption for 56 days of research on the treatment of *Tamarin-Pro* extract through drinking water did not show a significant difference (P>0.05) with control (D).

Table 1. Effects of supplementation Tamarin-Pro level on drinking water on egg production, feeds digestibility and N-NH₃ in laying hens.

Variables	Groups ¹				SEM ²
	A	B	C	D	
Initial body weight (g)	1670.35a ³⁾	1675.08a	1679.31a	1668.74a	19.052
Final body weight (g)	1716.92a	1762.07a	1708.94a	1754.63a	32.904
Egg mass (g/bird/8weeks)	3163,01ab	3183,51a	3217,4a	3033,80b	42,046
Egg production (egg/bird/8weeks)	49,98a	50,18a	50,42a	49,00a	0,641
Feed consumption (g/bird/8weeks)	7265,61a	6888,54a	6912,26a	6963,48a	109,82 5
Feed conversion ratio (feed consumption/egg mass)	2,29a ³⁾	2,14b	2,15b	2,29a	0,017
Dry matter digestibility (%)	75,04b	78,59a	78,92a	74,81b	0,252
Organic matter digestibility (%)	77,09b	80,77a	81,02a	76,9b	0,389
N-NH ₃ (m.Mol)	22,27b	21,66b	23,34b	42,47a	2,376

Note:

1. The drinking water treatments included: drinking water with 2% *Tamarin-Pro* extract (A); drinking water with 4% *Tamarin-Pro* extract (B); drinking water with 6% *Tamarin-Pro* extract (C); and drinking water as a control (D), respectively.
2. SEM: Standard Error of Treatment Means
3. Means with different superscripts within raw values are significantly different ($P < 0.05$)

The administration of *Tamarin-Pro* water extract on drinking water, namely in group B and C chickens, was significant ($P < 0.05$) higher each: 4.93% and 6.05% than controls (without *Tamarin-Pro* extract). The average value of the feed conversion ratio (FCR) in the control chicken group (D) was 2.29/head (Table 2). The average FCR value in group A chickens was 0.18% not significant ($P > 0.05$) lower than the control. However, the average FCR values in the chicken group B and C, were: 5.59% and 6.28% significantly different ($P < 0.05$) lower than the control (D).

Observations on the digestibility of feed dry matter turned out to increase significantly ($P < 0.05$) in the presence of 4% and 6% *Tamarin-Pro* extract in drinking water, ie 5.05% and 5.49% higher than controls. Similarly, the organic matter digestibility of feed in group B and C chickens was: 5.03% and 5.36% higher than the control (D).

The content of N-NH₃ in chicken excreta treated with *Tamarin-Pro* extract, namely chickens in Groups A, B, and C, were decreased significantly ($P < 0.05$) lower by 47.56%; 48.99%; and 45.04%, rather than controls or chicken in Group D.

DISCUSSION

The total number of eggs and feed consumption for 56 days of research on the treatment of fermented *Tamarindus indica* extract by yeast *Saccharomyces spp* (*Tamarin-Pro*) through drinking water did not show a significant difference with control (D). This is due to the fact that the metabolic energy content and dietary nutrients for the four treatments are the same. Reported by Bidura *et al.* (2017); Ekayuni *et al.* (2017) and Sanchez *et al.* (2005) which shows that herbal extracts do not have toxic effects or do not contain factors that limit intake which are opposite to absorption of nutrients.

The extract of fermented herbal products has not been able to increase egg production and feed consumption. According to Shivaramaiah *et al.* (2011), among these directly given microbes, their survival through digestion, growth in the digestive tract, and excretion through feces. *Tamarin-Pro* supplementation in drinking water has no significant effect on egg production. The same thing was reported by Davis and Anderson (2002) that there was no increase in egg production in laying hens added with probiotic bacteria, including *Lactobacillus* and *Bacillus*. This is also in accordance with the study of Kalavathy *et al.* (2009) who reported that probiotic supplementation had no significant effect on egg production in chickens. Egg production in chickens fed probiotic bacteria and yeast did not differ significantly from control chickens aged 40 to 52 weeks (Balevi *et al.*, 2001). Similarly, the results found by researchers, such as Ahmed *et al.* (2017) finding herbal leaf flour has no effect on feed consumption. Similar things Paguia *et al.* (2014) found that the supply of *Moringa* leaf flour did not affect the consumption of rations.

These conflicting results regarding growth performance responses to naturally fermented or extracted herbs are caused by different types of herbs, concentrations of herbs, and herbal processing methods (Windisch *et al.*, 2008; Embuscado, 2015). Decrease in feed consumption can be attributed to the presence of phytate, tannins, and saponins, in herbal leaves which all can reduce feed consumption, due to their unpleasant nature and binding to nutrients (Anhwange *et al.*, 2004).

The use of *Tamarindus* leaf extract fermented by probiotics (*Tamarin-Pro*) on the drinking water were significantly increased egg mass and feed efficiencies in group B and C chickens. *Tamarin-Pro* leaf water extract can be used as an effective feed supplement for laying poultry to improve feed efficiency (Akhouri *et al.*, 2013; Sanchez *et al.*, 2005). The main way of action of this active ingredient is the inhibition of microbial pathogens and endotoxins in the chicken intestine and increasing pancreatic activity, producing metabolism and better utilization of nutrients (Windisch *et al.*, 2008; Grashorn, 2010).

Digestion of dry matter and organic matter in hens groups C and D, was higher than the control. As recommended by Hernandez *et al.* (2004), that plant extract supplements can improve nutrient digestibility in the digestive tract of poultry. Herbal extracts (Garlic) can increase the activity of pancreatic enzymes and microenvironmental conditions for better utilization of nutrients in mice (Ramakrishna *et al.*, 2003).

Plants are rich in various secondary metabolites, such as tannins, terpenoids, alkaloids, and flavonoids, which have been found in vitro to have antimicrobial properties. These active compounds in the digestive tract of chickens can help absorb nutrients. As reported by Adibmoradi *et al.* (2006), that herbal active compounds in garlic can increase villous height and crypt depth, and reduce epithelial thickness and the number of villous cells in the duodenum, jejunum, and poultry ileum. Increased height of villi, as well as the thickness of epithelium and cup in the duodenum, jejunum and ileum will increase nutrient uptake (Nusairate, 2007). The results of the study by Bidura *et al.* (2017) found that the administration of *Sauropus* leaf extract in drinking water can significantly improve feed efficiency in laying hens.

The administration of *Tamarindus* leaf extract fermented by probiotics (*Tamarin-Pro*) in drinking water, namely in Group B and C hens, significantly reduced ammonia gas content in chicken excreta compared to Group D (control). *Tamarin-Pro* contains yeast *Saccharomyces spp* as a source of probiotics in drinking water drunk by chicken can reduce the concentration of ammonia gas in chicken excreta. The use of probiotic microbes in poultry is reported to be able to suppress urease enzyme activity and can reduce the amount of uric acid in the digestive tract of chickens, because uric acid has been used as a microbial protein (Chiang and Hsieh, 1995). According to Nguyen *et al.* (2019), the largest increase in growth performance, nutrient digestibility, faecal bacterial enumeration, and harmful gas emissions in weaned pigs was obtained when a probiotic mixture was added to feed as much as 0.3%. The same results were obtained when fermented probiotics in dry form were used as chicken daily feed (Lokman *et al.*, 2015).

It is assumed that feed fermentation generally improves the bacterial ecology of the digestive tract and the immune response in chicks, therefore, becomes a new model for future strategies to control chicken disease (Hasan *et al.*, 2016). Probiotics can maintain beneficial intestinal microflora, increase host resistance to enteric pathogenic bacteria, such as *Salmonella* and *Campylobacter species*, and produce a healthy digestive environment in poultry digestive

systems, feed conversion, and higher poultry performance (Vila *et al.*, 2009; Mountzoris *et al.*, 2010).

Supplementation of probiotics into rations significantly increases weight gain and decreases N-NH₃ levels in excreta of poultry (Chen *et al.*, 2009; Roni *et al.*, 2014; Puspani *et al.*, 2014). The same was reported by Bidura *et al.* (2014), that supplementation of *Saccharomyces* spp culture isolated from Bali cow feces by as much as 0.20% in diets can improve performance and reduce ammonia levels in broiler excreta.

CONCLUSION

We conclude that supplementation of fermented *Tamarindus indica* by *Saccharomyces* spp (*Tamarin-Pro*) extract on drinking water positively changed egg performance and feed digestibility of laying hens, but decreased N-NH₃ in hens excreta.

ACKNOWLEDGEMENTS

The authors would like to thank to Mr. Xanana Gusmao for the scholarship offered and Prof. Dr. Fransisco Martins as a Rector of National University, Timor Lorosa'e, East Timor for support, guidance and including research funding. The authors would like to thank to the Rector of Udayana University for their support during this study.

CONFLICT OF INTEREST DECLARATION

We certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

REFERENCES

- i. Abdelqader, A., Irshaid, R., and A.R. Al-Fataftah. 2013. Effects of dietary probiotic inclusion on performance, eggshell quality, cecal microflora composition, and tibia traits of laying hens in the late phase of production. *Trop. Anim. Health Prod.* 45, 1017-1024.
- ii. Adibmoradi, M., B. Navidshad, J. Seifdavati and M. Royan. 2006. Effect of dietary garlic meal on histological structure of small intestine in broiler chickens. *Jpn. Poult. Sci.* 43: 378-383.
- iii. Ahmed, S., Khaliq, A., Pasha, T.N., Mehmood, S., Hussain, K., Ahmad, S., Shaheen, M.S., Naeem, M and Shafiq, M. 2017. Effect of *Moringa oleifera* (Lam.) pods as feed additive on egg antioxidants, chemical composition and performance of commercial layers. *S. Afr. J. Anim Sci.* 47: 864-874.
- iv. Anhwange, B. A., Ajibola, V.O. and Oniye, S.J., 2004. Amino acid composition of the seeds of *Moringa oleifera* (Lam), *Detarium microcarpum* (Guill & Sperr) and *Bauhinia monandra* (Linn.). *J. Biol. Sci.* 9: 3-13
- v. AOAC. 2005. Official Methods of Analysis of AOAC International. 18th Ed. Association of Official Analytical Chemists, Arlington, USA.

-
- vi. Balevi, T, Ucan, US, Coskun, B, Kurtoglu, V, and I.S.Cetingul. 2001. Effect of dietary probiotic on performance and humoral immune response in layer hens. *Br. Poult. Sci.* 42, 456–461.
 - vii. Bidura, I.G.N.G., I.B.G. Partama, B.R.T. Putri, and N.L.Watiniasih. 2017. The effect of water extract of two leaves (*Allium sativum* and *Sauropus androgynus*) on the egg production and yolk cholesterol level in layer hens. *Pakistan Journal of Nutrition* Vol 15 (1): 23-31
 - viii. Bidura, IGNG., N. W. Siti and I.A. Putri Utami., 2014. Isolation of cellulolytic bacteria from rumen liquid of buffalo both as a probiotics properties and has CMC-ase activity to improve nutrient quality of soybean distillery by-product as feed. *International Journal of Pure & Applied Bioscience* Vol 2 (5): 10-18
 - ix. Bidura, I.G.N.G., N.W. Siti and I.B.G. Partama. 2019. Effect of probiotics, *Saccharomyces* spp.Kb-5 and Kb-8, in diets on growth performance and cholesterol levels in ducks. *South African Journal of Animal Science* Vol. 49 (2), 219-226
 - x. Bidura, I.G.N.G., Suyadnya, I.P., Mahardika, I.G., Partama, I.B.G., I.G.L.Oka, I.G.L. and I.G.A.I. Aryani. 2012. The implementation of *Saccharomyces spp.n-2* isolate culture (isolation from traditional yeast culture) for improving feed quality and performance of male Bali duckling. *Agricultural Science Research Journal* Vol. 2 (9), 486-492
 - xi. Carvalho, M.D.C.D. 2019. Supplementation Extract Fermented *Tamarindus Indica* by *Saccharomyces Spp* on Egg Production and Egg quality of Laying Hens. Disertation, Doctor Programs of Animal Science, Faculty of Animal Science, Udayana University, Denpasar, Indonesia
 - xii. Chen, K.L., Kho, W.L., You, S.H., Yeh, R.H., Tang, S.W. and C.W. Hsieh. 2009. Effects of *Bacillus subtilis* var. natto and *Saccharomyces cerevisiae* mixed fermented feed on the enhanced growth performance of broilers. *Poult. Sci.* 88, 309-315.
 - xiii. Chiang, S.H., and W.M. Hsieh. 1995. Effect of Direct Feed Microorganisms on Broiler Growth Performance and Litter Ammonia Level. *AJAS* 8 : 159-162
 - xiv. Davis, G.S. and K.E. Anderson. 2002. The effects of feeding the direct-fed microbial, primalac, on growth parameters and egg production in single comb white leghorn hens. *Poult. Sci.* 81, 755-759
 - xv. Ekayuni, A.A., I.G.N.G. Bidura, and I.B. Gaga Partama., 2017. 2017. The Effect of Water Extract of Two Leaves (*Moringa oleivera* and *Sauropus androgynus*) on Growth Performance and Meat Cholesterol Levels in Broilers. *J. Biol. Chem. Research.* Vol. 34, No. 1: 118-128
 - xvi. Embuscado, M.E., 2015. Spices and herbs: Natural sources of antioxidants – a mini review. *J. Func. Foods.* PP, 1-9.
 - xvii. Ezema, C. and Didacus C. Eze. 2015. Probiotic Effect of Yeast (*Saccharomyces cerevisiae*) on Hen-Day Egg Performance, Serum and Egg Cholesterol Levels in Laying Chicken. *Pakistan Journal of Nutrition* 14 (1): 44-46
 - xviii. Frankič T, Voljč M, Salobir J, Rezar V. 2009. Use of herbs and spices and their extracts in animal nutrition. *Acta Agric. Slov.* 94: 95-102.
-

-
- xix. Gheisar, M.M. and I.H. Kim (2018) Phytobiotics in poultry and swine nutrition – a review, Italian Journal of Animal Science, 17:1, 92-99, DOI: 10.1080/1828051X.2017.1350120
- xx. Grashorn, M.A., 2010. Use of phytobiotics in broiler nutrition: an alternative to infeed antibiotics. J. Anim. Feed. Sci. 19, 338-347.
- xxi. Hasan S.A. Jawad, I.H. Lokman, Saad A. Naji, A.B.Z. Zuki and A.B. Kassim. 2016. Effects of dietary supplementation of wet fermented feed with probiotic on the production performance of Akar Putra chicken. Asian J. Poult. Sci., 10: 72-77
- xxii. Hasanuddin, A., Rusdi and Rosmiaty Arief. 2017. Effects of Inclusion of Fermented Carrageenan By-products in the Basal Diet of Broiler Chickens on Growth Performance, Blood Profiles and Meat Composition. Int. J. Poult. Sci., 16 (5): 209-214)
- xxiii. Hernandez, F., Madrid, J., Garcia, V., Orengo, J. and Megías, M.D. 2004. Influence of two plant extracts on broilers performance, digestibility, and digestive organ size. Poultry Science, 83: 169-174.
- xxiv. Kalavathy, R., Abdullah N, Jalaludin S, Wong M, and Y.W. Ho. 2009. Effects of Lactobacillus cultures on performance of laying hens, and total cholesterol, lipid and fatty acid composition of egg yolk. J. Sci. Food Agric. 89, 482-486.
- xxv. Lei, K., Li, Y.L., Yu, D.Y., Rajput, I.R., and Li, W.F. 2013. Influence of dietary inclusion of Bacillus licheniformis on laying performance, egg quality, antioxidant enzyme activities, and intestinal barrier function of laying hens. Poultry Sci. 92, 2389-2395.
- xxvi. Lei, X.J., Hyeok Min Yun and In Ho Kim. 2018. Effects of dietary supplementation of natural and fermented herbs on growth performance, nutrient digestibility, blood parameters, meat quality and fatty acid composition in growing-finishing pigs , Italian Journal of Animal Science, 17:4, 984-993, DOI: 10.1080/1828051X.2018.1429955
- xxvii. Lokman, I.H., Jawad, S.H., Zuki, A.B.Z. and A.B. Kassim. 2015. Effect of dry probiotic supplemented fermented feed on production performance of akar putra chicken. Int. J. Poult. Sci.14, 420-426.
- xxviii. Mountzouris, K. C., P. Tsitsrikos, I. Palamidi, A. Arvaniti, M. Mohnl. G. Schatzmayr and K. Fegeros. 2010. Effect of probiotic inclusion levels in broiler nutrition on growth performance, nutrient digestibility, plasma immunoglobulins, and cecal microflora composition. Poult. Sci. 89: 58-67.
- xxix. Nusairat, B. M., 2007. Dietary supplementation of garlic (*Allium Sativum*): Influence on performance parameters, meat quality and humoral immune response in broiler chicks. M.S. Thesis, Jordan University of Science and Technology, Irbid, Jordan.
- xxx. Nguyen, D.H., C. M. Nyachoti and I. H. Kim. 2019. Evaluation of effect of probiotics mixture supplementation on growth performance, nutrient digestibility, faecal bacterial enumeration, and noxious gas emission in weaning pigs, Italian Journal of Animal Science, 18:1, 466-473, DOI: 10.1080/1828051X.2018.1537726
-

-
- xxx. Paguia, H.M., Paguia, R.Q., Balba, C. and Flores, R.C., 2014. Utilization and evaluation of *Moringa oleifera* L. as poultry feeds. 4th International Conference on Agriculture and Animal Science. 8, 343-7
- xxxii. Phuoc, T.L. and U. Jamikorn. 2017. Effects of probiotic supplement (*Bacillus subtilis* and *Lactobacillus acidophilus*) on feed efficiency, growth performance, and microbial population of weaning rabbits. *Asian-Australas J Anim Sci* 30:198-205
- xxxiii. Puspani, E., IGNG. Bidura, DPMA. Candrawati and N.W. Siti. 2014. Pollard in diets supplemented with yeast on broiler performance and ammonia-N concentration of excreta. *Journal of Biological and Chemical Research* 31 (2): 1048-1055
- xxxiv. Ramakrishna, R. R., K. Platel and K. Srinivasan. 2003. *In vitro* influence of spices and spice-active principles on digestive enzymes of rat pancreas and small intestine. *Nahrung* 47: 408-412.
- xxxv. Roni, N.G.K., E. Puspani, and I G.N.G. Bidura. 2014. Probiotics feeding management as an effort for pressuring body fat and excrete ammonia concentration of Bali drake. *Majalah Ilmiah Peternakan* 18 (3): 119-124
- xxxvi. Sanchez, N.R., E. Spornly and I. Ledin. 2005. Effect of different levels of foliage of *Moringa oleifera* to creole dairy cows in intake, digestibility, milk production and composition. *Livest., Sci.* 2810: 8
- xxxvii. Shivaramaiah S, Pumford NR, Morgan MJ, Wolfenden RE, Wolfenden AD, Torres-Rodríguez A, Hargis BM, and Tellez G. 2011. Evaluation of *Bacillus* species as potential candidates for direct-fed microbials in commercial poultry. *Poult. Sci.* 90, 1574-1580.
- xxxviii. Tufarelli, V., A. M. Crovace, G. Rossi and V. Laudadio. 2017. Effect of a dietary probiotic blend on performance, blood characteristics, meat quality and faecal microbial shedding in growing-finishing pigs. *South African Journal of Animal Science*, 47 (6): 875-882
- xxxix. Vila, B., A. Fontgibell, I. Badiola, E. Esteve-Garcia, G. Jiménez, M. Castillo and J. Brufau. 2009. Reduction of *Salmonella entericav.* Enteritidis colonization and invasion by *Bacillus cereus* var. *toyoi* inclusion in poultry feeds. *Poult. Sci.* 88:975-979.
- xl. Windisch, W., Schedle, K., Plitzner, C. and Kroismayr, A. 2008. Use of phytogetic products as feed additives for swine and poultry. *J. Anim. Sci.* 86, 140-148.
- xli. Zhen, Y.G., W. Zhao, X. Chen, L.J. Li, H.G. Lee, X.F. Zhang and T. Wang. 2019. Effects of yeast culture on broiler growth performance, nutrient digestibility and caecal microbiota. *S. Afr. J. Anim. Sci.* vol. 49 (1): 99-108